

FLUOPYRAM	GROUP	7	FUNGICIDE
TRIFLOXYSTROBIN	GROUP	11	FUNGICIDE



## FIFRA Section 2 (ee) Recommendation

### FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NEW YORK

#### BROADFORM®

#### For Control of Beech Leaf Disease on Beech Trees with Foliar Application

EPA Reg. No. 432-1537

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Plant	Target Disease	Application Rate	Interval between applications (days)
Beech Tree	Beech Leaf Disease (BLD)	8 fluid ounces/100 gallons	7 – 14 until the threat of disease is over
	<ul style="list-style-type: none"> <li>Do not apply more than 27.3 fl. oz. of BROADFORM per acre per year.</li> <li>Applications are to be made through hydraulic sprayers operating between 150-350 psi.</li> <li>Use of a spreader sticker is recommended to improve efficacy for BLD.</li> </ul>		

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

**For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).**

*As with any crop-protection product, always read and follow label directions.*

*For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

**Bayer Environmental Science**  
A Division of Bayer CropScience LP  
5000 CentreGreen Way, Suite 400  
Cary, NC 27513